



PNEUMOCOCCAL POLYSACCHARIDES VACCINE

PNEUMOVAX 23®

WHAT YOU NEED TO KNOW



1. Pneumococcal disease

Pneumococcal disease is caused by *Streptococcus pneumoniae* bacteria. It is a leading cause of vaccine-preventable illness and death in the United States and Europe. Anyone can get pneumococcal disease, but some people are at greater risk than others:

- People 65 years and older.
- The very young.
- People with certain health problems.
- People with a weakened immune system.
- Smokers.

Pneumococcal disease can lead to serious infections of the:

- Lungs (pneumonia)
- Blood (bacteremia)
- Covering of the brain (meningitis).

Pneumococcal pneumonia kills about 1 out of 20 people who get it. Bacteremia kills about 1 person in 5, and meningitis about 3 people in 10. People with the health problems described in Section 3 of this statement may be more likely to die from the disease.

2. What is Pneumovax 23® vaccine?

- Treatment of pneumococcal infections with penicillin and other drugs used to be more effective. But some strains of the disease have become resistant to these drugs. This makes prevention of the disease, through vaccination, even more important.
- Pneumococcal polysaccharide vaccine (PPSV) / Pneumovax 23® protects against 23 types of pneumococcal bacteria, including those most likely to cause serious disease.
- Most healthy adults who get the vaccine develop protection to most or all of these types within 2 to 3 weeks of getting the shot. Very old people, children under 2 years of age, and people with some long-term illnesses might not respond as well, or at all.

Another type of pneumococcal vaccine (pneumococcal conjugate vaccine, or PCV) is routinely recommended for children younger than 2 years of age.

3. Who should get Pneumovax 23® vaccine?

- All adults 65 years of age and older.
- Anyone 2 through 64 years of age who has a long-term health problem such as:
 - ⇒ heart and or a lung disease
 - ⇒ sickle cell disease
 - ⇒ diabetes
 - ⇒ alcoholism
 - ⇒ cirrhosis
 - ⇒ leaks of cerebrospinal fluid or cochlear implant
- Anyone 2 through 64 years of age who has a disease or condition or is taking a drug or treatment that lowers the body's resistance to infection
- Any adult 19 through 64 years of age who is a smoker or has asthma

Pneumovax 23® may be less effective for some people, especially those with lower

resistance to infection. But these people should still be vaccinated, because they are more likely to have serious complications if they get pneumococcal disease.

Children who often get ear infections, sinus infections, or other upper respiratory diseases, but who are otherwise healthy, do not need to get Pneumovax 23 vaccine because it is not effective against those conditions.

4. How many doses of Pneumovax 23 vaccine are needed, and when?

Usually only one dose of PPSV is needed, but under some circumstances a second dose may be given.

- A second dose is recommended for people 65 years and older who got their first dose when they were younger than 65 and it has been 5 or more years since the first dose.

A second dose is recommended for people 2 through 64 years of age who:

- have a damaged spleen or no spleen.
- have sickle-cell disease.
- have HIV infection or AIDS.
- have cancer, leukemia, lymphoma, multiple myeloma.
- have had an organ or bone marrow transplant.
- are taking medication that lowers immunity (such as chemotherapy or long-term steroids)

When a second dose is given, it should be given 5 years after the first dose.

5. Some people should not get Pneumovax 23 vaccine or should wait:

- Anyone who has had a life-threatening allergic reaction to PPSV should not get another dose.
- Anyone who has a severe allergy to any component of a vaccine should not get that vaccine. Tell your provider if you have any severe allergies.
- Anyone who is moderately or severely ill when the shot is scheduled may be asked to wait until they recover before getting the vaccine. Someone with a mild illness can usually be vaccinated.
- While there is no evidence that PPSV is harmful to either a pregnant woman or to her fetus, as a precaution, women with conditions that put them at risk for pneumococcal disease should be vaccinated before becoming pregnant, if possible.

5. What are the risks from Influenza vaccine?

A vaccine, like any medicine, could possibly cause serious problems such as severe allergic reactions. ***The risk of Influenza vaccine causing serious harm, or death, is extremely small.***

- About half of people who get PPSV have mild side effects, such as redness or pain where the shot is given.
- Less than 1% develops a fever, muscle aches, or more severe local reactions.

If these reactions occur, they usually last for 1 or 2 days.

6. What if there is a moderate or severe reaction?

What should I look for ?

- Any unusual condition, such as a high fever or behavior changes. Signs of a serious allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do ?

- Call a doctor, or get the person to a doctor right away.
- Tell your doctor what happened, the date and time it happened, and when the vaccination was given.

7. How can I learn more?

- Ask your doctor or nurse. They can give you the vaccine package insert or suggest other sources of information.
- Contact the American Centers for Disease Control and Prevention (CDC):
Visit CDC websites at: www.cdc.gov/nip